

# Evaluation of Hyperuricaemia

Suggested scheme for evaluation of Hyperuricaemia

---

## Exclude

---

### Potentially correctable contributory factors

- Obesity
- Alcohol
- Hypertriglyceridaemia
- Drugs (especially thiazides, but also diuretics, low doses of salicylates, nicotinic acid, pyrazinamide, ethambutol, and cyclosporin)
- Hypertension
- Low fluid intake

---

## Consider

---

### High purine intake

Diet (meats, yeast products)

---

### Increased urate production

#### Primary

- Idiopathic
- Enzyme defects

#### Secondary

- Blood dyscrasias
- Infectious mononucleosis
- Malignancy
- Cytotoxic therapy
- Psoriasis
- Alcoholism
- Prolonged exercise

---

### Decreased renal excretion

#### Primary

- Syndrome-X (insulin resistance, dyslipidaemia [increased TG, low HDL-cho], obesity, hypertension, hyperuricaemia)
- Idiopathic

#### Secondary

- Renal failure. If renal failure is causing the hyperuricaemia, serum creatinine will be  $> 0.40$  mmol/L, serum urate will be  $< 0.65$  mmol/L, and the ratio of urine urate to creatinine will be  $< 0.7$ . If hyperuricaemia is causing the renal failure, then serum urate will be  $> 0.7$  mmol/L and the ratio of urine urate to creatinine will be  $> 0.7$ .
- Dehydration
- Diuretics
- Ketonaemia (starvation, diabetes mellitus)
- Hyperlactataemia (alcohol, toxaemia of pregnancy)
- Drugs
- Hyperparathyroidism

---

## If the cause is obscure

Consider 24-hour urinary excretion rate before and after a 5-day low-purine diet:

	<b>Normal diet</b> (mmol/24hr)	<b>Low-purine diet</b> (mmol/24hr)
High purine intake	$> 6.0$	$< 4.0$
Increased urate production	$> 6.0$	$> 4.5$
Decreased renal excretion	$< 6.0$	$< 4.0$ (often $< 2.0$ )