

Evaluation of Lactate Dehydrogenase Isoenzymes (LDI)

Increased Lactate Dehydrogenase (LDH)

Exclude *in vitro* haemolysis

Perform Isoenzymes (if cause of LDH elevation is unclear)

Elevation of LD1

Myocardial infarction

Red blood cell diseases (e.g. haemolytic anaemia), B12 deficiency (↑ MCV)

In vitro haemolysis

Some muscle dystrophies (e.g. Duchenne muscular dystrophy)

Kidney disease

Kidney transplant rejection

Testicular/Germ cell tumours/rare neuroendocrine tumours

Intense exercise training

Elevation of LD2

Infections

Lung diseases (LD 2 & 3)

Congestive heart failure

Lymphocyte turnover (e.g. lymphomas/EBV) (LD 2 & 3)

Elevation of LD3

Skin (psoriasis)

Lung disease/injury

Lymphocyte turnover (e.g. lymphomas/EBV)

Splenic disorders (infarct)

Platelet destruction

Elevation of LD4

Placenta disorders

Elevation of LD5

Liver diseases: hepatitis, toxins (paracetamol)

Skeletal muscle diseases/injuries

Some intestinal problems

Pleural fluid: neutrophils (LD 4 & 5)

Elevation of multiple LDIs

Elevation of LD1, LD2, and LD5 may be caused by strenuous exercise.

Elevation of all the LDIs may be caused by injury to multiple organs (e.g. congestive heart failure, advanced cancers, autoimmune diseases, or shock).

Increases in Midzone Levels

Malignant pattern (crescendo-decrescendo pattern)

Storage

Marked LD4/LD5 decrease occurs if sample stored below room temperature. If liver/skeletal muscle LDI is suspected, a fresh specimen is recommended.

Extra isoenzyme bands

Macrocomplex (usually benign)
